

EVALUATION OF THE EFFECT OF REGULATED DEFICIT IRRIGATION ON VITIS VINIFERA CABERNET SAUVIGNON PHYSIOLOGICAL TRAITS AND FINAL FRUIT COMPOSITION

Authors: Sebastian VARGAS^{1*}, Francesco GUIDI^{1,2}, Edmundo BORDEU³, Alvaro GONZALEZ¹, Samuel ORTEGA-FARÍAS⁴

¹Centro de Investigación e Innovación de Viña Concha y Toro, Ruta K-650 km 10 Penciahue, Chile

²École Supérieure d'Agriculture d'Angers, 55 rue Rabelais 49007 ANGERS, France

³Pontificia Universidad Católica de Chile, Av. Vicuña Mackenna 4860, Santiago, Chile

⁴Centro de Investigación y Transferencia en Riego y Agroclimatología (CITRA), Universidad de Talca, Av. Lircay s/n, Talca, Chile

*Corresponding author: sebastian.vargass@conchaytoro.cl

Abstract:

Context and purpose of the study - Climate change establishes challenges, as well as opportunities for many sectors, and markedly the wine sector. The wine industry plays a weighty role in Chile's economy, where contributes considerably to the country financial development. Counting the vulnerability of the country to such changes, it is fundamental to enact preventive measures aiming at besting the resources management, above all water necessities for cultivated crops. Optimal irrigation in grapevines could be accomplished by means of physiological data based programming and final grape and wine chemical and sensory performance. This study aims to understand the impacts of different levels of deficit irrigation on a large amount of chemical markers from aroma to non-volatile compounds, and the final impact on sensory profile.

Material and methods - A regulated deficit irrigation (RDI) experimental trial that was conducted in a commercial vineyard of Cabernet Sauvignon in the Maule valley in central Chile. Four regulated deficit irrigation (RDI) regimes were employed in four replicated blocks to replenish different portions of evapotranspiration (ET) from pea-size stadium until harvest. These managements were conceived 100% ET, 70% ET, 50-100% ET (50% ET before veraison and 100% ET afterward) and 25-100% ET (25% ET before veraison and 100% ET afterward). The following parameters were measured: midday stem water potential (Ψ_{stem}), stomatal conductance (gs), vine and grapes growth, yield, quality of must. GSMS for norisoprenoid, terpene, C6s compounds and methoxypyrazines concentration and HPLC for anthocyanin and low molecular weight phenols was used.

Results - Of definite interest were the outcomes from the grapes' evolution monitoring, as we had expectations of an alteration in their development in RDI conditions. Apart from the mere berry size, which showed significant differences between the treatments and control, no other variations have been registered. Surprisingly, the °Brix degrees were very alike, indicating that a reduced water availability does not always imply a faster maturation of the sugars in the grapes, as opposed to previous investigations that shows that a conventional irrigation may imply a delay in sugar accumulation. These circumstances additionally, allowed us to harvest all of the four regimes at the same time, thus providing optimal comparison bases. Significant differences were found in several traits, from lower concentration of malic acid in RDI treatments to higher concentrations of anthocyanin and some specific flavonoids like quercetin and miricetins. In the case of aroma compounds, our partial results indicate a significant effect of the RDI in increasing the concentration of 3-Isobutyl-2-methoxypyrazine. This results might be linked to increased light interception in the RDI treatments, who tend to defoliate early in the season.

Keywords: regulated deficit irrigation, Cabernet Sauvignon, stem water potential, stomatal conductance, yield components, IBMP, Low molecular weight phenols.

1. Introduction.



CENTRO DE
INVESTIGACIÓN
E INNOVACIÓN
VIÑA CONCHA Y TORO

EVALUATION OF THE EFFECT OF REGULATED DEFICIT IRRIGATION ON VITIS VINIFERA CABERNET SAUVIGNON PHYSIOLOGICAL TRAITS AND FINAL FRUIT COMPOSITION



Sebastian VARGAS^{1*}, Francesco GUIDI^{1,2}, Edmundo BORDEU³, Alvaro GONZALEZ¹, Samuel ORTEGA-FARIAS⁴

¹Center from Research and Innovation, Viña Concha y Toro, Chile

²École Supérieure d'Agriculture d'Angers, France

³Pontificia Universidad Católica de Chile, Santiago, Chile

⁴Centro de Investigación y Transferencia en Riego y Agroclimatología (CITRA), Universidad de Talca, Chile

INTRODUCTION

Water availability in agriculture has become a limiting factor, mainly because of constant drought associated with the Niña phenomenon which leads to fewer precipitations in Chile.

Optimal irrigation in grapevines could be accomplished by means of physiological data based programming and final grape and wine chemical and sensory performance. This study aims to understand the impacts of different levels of deficit irrigation on a large amount of chemical markers from aroma to non-volatile compounds, and the final impact on sensory profile.

MATERIALS & METHODS

A regulated deficit irrigation (RDI) experimental trial that was conducted in a commercial vineyard of Cabernet Sauvignon in the Maule valley in central Chile. Four regulated deficit irrigation (RDI) regimes were employed in four replicated blocks to replenish different portions of evapotranspiration (ET) from pea-size stadium until harvest. These managements were conceived X0: 100% ET, X1: 70% ET, X2: 50-100% ET (50% ET before veraison and 100% ET afterward) and X3: 25-100% ET (25% ET before veraison and 100% ET afterward). The following parameters were measured: midday stem water potential (Ψ_{stem}), stomatal conductance (gs), vine and grapes growth, yield, quality of must. GSMS for norisoprenoid, terpene, C6s compounds and methoxypyrazines concentration and HPLC for anthocyanin and low molecular weight phenols was used.




Figure 1. Midday stem water potential of treatments during the season. Different letters in vertical order indicate significant differences $p \leq 0,05$.




Figure 2. Principal components analysis of sensory attributes of wines of plant of the different treatments.

RESULTS

RDI technique reduces the sources, the foliage in this case expressed as pruning weight but also the Sinks, reducing the total yield of grapes produced, mainly by a lighter clusters with fewer and smaller berries (Table 1, Figure 5). Since sources and sinks are reduced rava index is not affected.

Table 1. Yield components by treatment. Different letters in vertical order indicate significant differences $p \leq 0,05$.

Treatment	Yield/ha (kg)	Pruned/Plant (kg)	Rava Index	Berry average weight (g)	Cluster weight (g)	Foliar weight (g)	Berries number
X0	16154 ^a	1.03 ^a	4,41	1.10 ^a	173 ^a	9.82 ^a	147 ^a
X1	15494 ^a	0.86 ^b	5,13	1.00 ^b	146 ^b	8.36 ^b	136 ^{ab}
X2	13,474 ^{ab}	0.90 ^b	4,11	0.85 ^b	114 ^b	7.81 ^{bc}	123 ^b
X3	11,982 ^b	0.90 ^b	3,73	0.82 ^b	105 ^b	6.77 ^c	120 ^b

This stable condition between sources and sinks might be the responsible of not having differences in grape sugar accumulation (Table 2). But in general terms an increase in the concentration of quality associated chemical markers is observed with reduced irrigation, higher concentrations of polyphenols like anthocyanins and some flavonoids, quercetin and miricetins mainly.

Table 2. Non volatile chemical characterization of grapes by treatment. Different letters in vertical order indicate significant differences $p \leq 0,05$.

Treatment	Brix	Malic acid (g)	pH	Total acidity	Total polyphenols (mg/L)	Quercetin (mg/L)	Resveratrol (mg/L)	Ac. Malic (mg/L)	Flavonoids (mg/L)	TPM (mg)
X0	24,1	1,67 ^a	3,53	2,91	707 ^b	78,0 ^b	94,39	4,60	136,9 ^b	235,9
X1	24,3	1,52 ^{ab}	3,53	2,85	688 ^b	84,7 ^{ab}	68,08	5,73	142,4 ^b	216,2
X2	24,2	1,28 ^{bc}	3,49	2,92	755 ^{ab}	97,1 ^a	73,69	8,35	181,0 ^a	263,0
X3	24,1	1,06 ^c	3,53	3,00	905 ^a	100,2 ^a	59,50	8,08	183,9 ^a	251,5

The experiment only shows an increased concentration of anthocyanins (Figure 3), but not effect on the concentration expressed by mg per Berry. So no prove of an effect on the synthesis or degradation of this molecules.

In the other hand, and increased concentration of 3-isobutyl-2-methoxypyrazine (IBMP) by litter and by Berry was possible to observe, particular for the 70% Etc treatment (Figure 4). So either a lower degradation or a higher synthesis.




Figure 3. Concentration of different anthocyanins in grape samples. Different letters in horizontal order indicate significant differences, for particular anthocyanin inside a bar, or for the total anthocyanin on the top. $p \leq 0,05$.

CONCLUSIONS

Of definite interest were the outcomes from the grapes' evolution monitoring, as we had expectations of an alteration in their development in RDI conditions. Apart from the mere berry size, which showed significant differences between the treatments and control, no other variations have been registered. Surprisingly, the °Brix degrees were very alike, indicating that a reduced water availability does not always imply a faster maturation of the sugars in the grapes, as opposed to previous investigations that shows that a conventional irrigation may imply a delay in sugar accumulation. These circumstances additionally, allowed us to harvest all of the four regimes at the same time, thus providing optimal comparison bases. Significant differences were found in several traits, from lower concentration of malic acid in RDI treatments to higher concentrations of anthocyanin and some specific flavonoids like quercetin and miricetins. In the case of aroma compounds, our partial results indicate a significant effect of the RDI in increasing the concentration of 3-Isobutyl-2-methoxypyrazine. This results might be linked to increased light interception in the RDI treatments, who tend to defoliate early in the season.




Figure 4. Concentration by litter and by berry of 3-isobutyl-2-methoxypyrazine (IBMP) in grape samples. Different letters in horizontal order indicate significant differences between treatments. $p \leq 0,05$.



Figure 5. Cluster of the 4 different treatments, from left to right, Control 100% ET, 70% ET, 50-100% ET, 25-100% ET.

ACKNOWLEDGMENTS

This study is part of a research project accredited under R&D Chilean law by CORFO. We thank to all the Agricultural Management department of Viña Concha y Toro. Special Thanks to the CITRA team from Universidad de Talca and Riego OLIVOS SPA. for their enormous contribution.